

Is a Quantum Logic a Logic?

by **R. J. Greechie**¹⁾ and **S. P. Gudder**

Kansas State University, Dept. of Mathematics, Manhattan, Kansas, USA
and University of Denver, Denver, Colorado, USA

(1. V. 70)

In a recent study Jauch and Piron [2] have considered the possibility that a quantum proposition system is an infinite valued logic. They argue that if this is the case then for any two propositions p and q there must exist a conditional proposition $p \rightarrow q$. Following Lukasiewicz [3] the truth value $[p \rightarrow q]$ of the conditional $p \rightarrow q$ is defined as follows: $[p \rightarrow q] = \min\{1, 1 - [p] + [q]\}$ where $[p]$ and $[q]$ are the truth values of p and q respectively. Here $[p] = 1$ is interpreted as ' p is true'. Note that $[p] = 1$ and $[p \rightarrow q] = 1$ implies $[q] = 1$ so we have a law of deduction, which is a property that any reasonable logic should possess. Notice further that if $[p \rightarrow q] = 1$ and $[q \rightarrow r] = 1$ then $[p \rightarrow r] = 1$ so that implication is transitive as it should be.

Let \mathcal{L} be an orthomodular poset (representing some quantum proposition system) and let \mathcal{S} be an order determining (full in [1]) set of states on \mathcal{L} . We further assume that if $m_1, m_2 \in \mathcal{S}$, then $1/2 m_1 + 1/2 m_2 \in \mathcal{S}$, that is, \mathcal{S} is closed under the formation of mid-points. We say that $a, b \in \mathcal{L}$ are *conditional* if there exists $c \in \mathcal{L}$ such that for all $m \in \mathcal{S}$ $m(c) = \min\{1, m(a') + m(b)\}$. If c exists it is unique. We call c the *conditional* of a and b and write $c = a \rightarrow b$. We say that \mathcal{L} (or, more correctly, the pair $(\mathcal{L}, \mathcal{S})$) is *conditional* if every pair $a, b \in \mathcal{L}$ are conditional. Now if \mathcal{L} is to be a logic with a law of deduction then \mathcal{L} must be conditional. Jauch and Piron [2] have shown that standard proposition systems (that is, ones that are isomorphic to the lattice of all closed subspaces of a Hilbert space) are not conditional and thus cannot be logics in the usual sense. We generalize their results to the orthomodular posets \mathcal{L} considered above. In fact we obtain the strong result that \mathcal{L} is conditional if and only if $\mathcal{L} = \{0, 1\}$. We then characterize the pairs $a, b \in \mathcal{L}$ which are conditional.

Undefined terms appear in [1]. If $a \leq b'$ we write $a + b$ for $a \vee b$. If $a \leq b$ we write $b - a$ for $b \wedge a'$. We first state a useful lemma whose simple proof is left to the reader.

Lemma 1. (i) $m(a \rightarrow b) = 1$ if and only if $m(a) \leq m(b)$; $m(a \rightarrow b) = m(a') + m(b)$ if and only if $m(b) \leq m(a) = 1$.

(ii) $m(a \rightarrow b) = m(b)$ if and only if $m(b) = 1$ or $m(a) = 1$.

This lemma will be frequently used without further comment.

Theorem 2. \mathcal{L} is conditional if and only if $\mathcal{L} = \{0, 1\}$.

¹⁾ Research supported in part by the National Science Foundation Grant Number GP-11241 and in part by the Fonds National Suisse.

Proof: Clearly $\{0, 1\}$ is conditional; in fact $1 = 0 \rightarrow 1$ and $0 = 1 \rightarrow 0$. Now let \mathcal{L} be conditional and suppose there exists $a \in \mathcal{L} - \{0, 1\}$. Then $c = a \rightarrow a'$ exists and $m(c) = \min \{1, 2 m(a')\}$. Since \mathcal{S} is order determining $a' \leq c$. Hence there exists $b \in \mathcal{L}$ such that $a' + b = c$. Now $m(b) = m(c) - m(a') = \min \{m(a), m(a')\}$. Thus $m(b) \leq 1/2$ for all $m \in \mathcal{S}$. It follows that $b \leq b'$ since \mathcal{S} is order determining. Hence $b = 0$ and $c = a'$. Thus $m(c) = \min \{1, 2 m(c)\}$ and hence $m(c) = 0$ or 1 for all $m \in \mathcal{S}$. Moreover since $0 < c < 1$ there exist $m_1, m_2 \in \mathcal{S}$ with $m_1(c) = 0$ and $m_2(c) = 1$. Letting $m = 1/2 m_1 + 1/2 m_2$ we have $m(c) = 1/2$, a contradiction. Hence $\mathcal{L} = \{0, 1\}$.

We have seen that, for non-trivial posets \mathcal{L} , not every pair of elements is conditional. We now study the properties of pairs of elements that are conditional.

Lemma 3. If $a \rightarrow b$ and $a' \vee b$ exist and are equal then $a C b$.

Proof: There exists $d \in \mathcal{L}$ such that $b + d = a' \vee b$. We show $d \leq a'$. Otherwise there exists $m \in \mathcal{S}$ such that $m(d) > m(a')$. Then $m(a' \vee b) = m(b) + m(d) > 1 - m(a) + m(b)$ so $m(a) > m(b)$. Hence $m(a' \vee b) = m(a \rightarrow b) = 1 - m(a) + m(b)$, a contradiction. Now there exists $e \in \mathcal{L}$ with $d + e = a'$. We show $e \leq b$. Otherwise there exists $m \in \mathcal{S}$ with $m(e) > m(b)$. Then $m(a') = m(d) + m(e) > m(d) + m(b) = m(a' \vee b) \geq m(a')$, a contradiction. Hence there exists $f \in \mathcal{L}$ with $b = f + e$, $a' = d + e$ and $f \leq b \leq d'$ so that $a' C b$. Thus $a C b$.

Lemma 4. If $c = a \rightarrow b$ exists then $a' \leq c$ and $b \leq c$.

Proof: If $a' \leq c$ then there exists $m \in \mathcal{S}$ such that $m(c) < m(a')$. Hence $m(c) < 1$ and $1 - m(a) + m(b) = m(c) < 1 - m(a)$. Thus $m(b) < 0$, a contradiction. That $b \leq c$ is immediate.

We say that \mathcal{S} is *sufficient* if $0 \neq a \in \mathcal{L}$ implies there exists $m \in \mathcal{S}$ with $m(a) = 1$.

Theorem 5. Let \mathcal{S} be sufficient and assume that $a' \vee b$ exists. Then $a \rightarrow b$ exists if and only if $a \leq b$ or $b \leq a$.

Proof: Clearly, if $a \leq b$ then $a \rightarrow b = 1$ and if $b \leq a$, then $a \rightarrow b = a' + b$. Conversely, assume $c = a \rightarrow b$ exists. By Lemma 4 $c \geq a' \vee b$. Hence there exists $d \in \mathcal{L}$ such that $(a' \vee b) + d = c$. Suppose $d \neq 0$. Then there exists $m \in \mathcal{S}$ such that $m(d) = 1$. Hence $m(a') = m(b) = 0$ and $m(c) = 1 - m(a) + m(b) = 0$, a contradiction. Therefore $d = 0$ and $c = a' \vee b$. It now follows from Lemma 3 that $a C b$. Suppose a and b are not comparable. Then $a \wedge b < a$ and $a \wedge b < b$. Hence there exists $m_1, m_2 \in \mathcal{S}$ such that $m_1(a - (a \wedge b)) = 1$ and $m_2(b - (a \wedge b)) = 1$. It follows that $m_1(a) = m_2(b) = 1$ and $m_1(b) = m_1(a \wedge b) = m_2(a) = m_2(a \wedge b) = 0$. Let $m = 1/2 (1/2 m_1 + 1/2 m_2) + 1/2 m_1 = 3/4 m_1 + 1/4 m_2$. Then $m(a \wedge b) = 0$ and $m(b) = 1/4 < 3/4 = m(a)$. Hence $m(a') + m(b) = m(c) = m(a' \vee b) = m(a' + (a \wedge b)) = m(a') + m(a \wedge b)$. Thus $m(b) = m(a \wedge b)$, a contradiction.

Corollary 6. Let \mathcal{S} be sufficient and $a' \vee b$ exist. If $a \rightarrow b$ exists, then $a \rightarrow b = a' \vee b$, $b \rightarrow a$ exists, $b' \vee a$ exists, and $b \rightarrow a = b' \vee a$.

The proofs of the previous theorems depend heavily on the fact that \mathcal{S} is order determining, sufficient or both. If we strengthen \mathcal{S} still further we obtain a stronger result. We say that \mathcal{S} is *strongly order determining* if $\{m \in \mathcal{S}: m(a) = 1\} \subset \{m \in \mathcal{S}: m(b) = 1\}$

implies that $a \leq b$. It can be shown that strongly order determining implies both order determining and sufficiency. (The converse fails; see [1].) Notice that the set of states on the lattice of all closed subspaces of a Hilbert space is strongly order determining.

Theorem 7. If \mathcal{S} is strongly order determining, then $a \rightarrow b$ exists if and only if $a \leq b$ or $b \leq a$.

Proof: As in Theorem 5, if a and b are comparable, then $a \rightarrow b$ exists. Now assume $c = a \rightarrow b$ exists. Suppose $a \not\leq b$ and $b \not\leq a$. Then there exists $m_0, m_1 \in \mathcal{S}$ such that $m_0(a) = 1$, $m_0(b) < 1$, $m_1(a) < 1$ and $m_1(b) = 1$. Note that $m_0(c) = m_0(b)$ and $m_1(c) = 1$. Let $m = 1/2 m_0 + 1/2 m_1$. Then $m(a) = 1/2 + 1/2 m_1(a) < 1$, $m(b) = 1/2 m_0(b) + 1/2 < 1$ and $m(c) = m(b)$. This last sentence contradicts Lemma 1 (ii). Hence a and b are comparable.

REFERENCES

- [1] R. J. GREECHIE, *An Orthomodular Poset with a Full Set of States not Embeddable in Hilbert Space*, Carib. J. Sci. Math. 7, 15 (1969).
- [2] J. M. JAUCH and C. PIRON, *What is 'Quantum-Logic'? quanta*, edited by P. G. O. FREUND, C. J. GOEBEL and Y. NAMBU (The University of Chicago Press, Chicago and London 1970), p. 166-181.
- [3] J. LUKASIEWICZ and A. TARSKI, *Investigations into the Sentential Calculus, Logic, Semantics and Metamathematics* (Clarendon Press, Oxford 1956).